

INVITED REVIEW SERIES:
RESPIRATORY HEALTH ISSUES IN THE ASIA-PACIFIC REGION

The effects of bushfire smoke on respiratory health

MARTINE DENNEKAMP AND MICHAEL J. ABRAMSON

Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Victoria, Australia

ABSTRACT

Bushfire smoke has the potential to affect millions of people and is therefore a major public health problem. The air pollutant that increases most significantly as a result of bushfire smoke is particulate matter (PM). During bushfire smoke episodes, PM concentrations are usually much higher than urban background concentrations, at which effects on respiratory health have been observed. The smoke can cover large areas including major cities and even small increases in the risk of respiratory health effects can cause large public health problems. The association between respiratory morbidity and exposure to bushfire smoke is consistent with the associations found with urban air pollution. Although using different methods, all studies looking at Emergency Department presentations in relation to a bushfire smoke event have found associations and most studies have also found an association with hospital admissions. However, only a few studies have distinguished between the effects of bushfire PM₁₀ (particles with a median aerodynamic diameter less than 10 µm) and background PM₁₀. These studies suggest that PM₁₀ from bushfire smoke is at least as toxic as urban PM₁₀, but more research is needed.

Key words: bushfire, forest fire, respiratory health, wildfire.

The Authors: Martine Dennekamp (MSc, PhD) is an epidemiologist with a particular interest in the area of air pollution and climate change on human health. Michael Abramson (MB BS, PhD, FRACP) is a respiratory physician and epidemiologist with broad ranging research interests in the causes and management of chronic respiratory diseases.

Correspondence: Martine Dennekamp, Centre for Occupational and Environmental Health, Department of Epidemiology and Preventive Medicine, Monash University, The Alfred Hospital, Commercial Road, Melbourne, Vic. 3004, Australia. Email: martine.dennekamp@monash.edu

Received 7 June 2010; invited to revise 29 June 2010, 23 August 2010; revised 2 August 2010, 15 September 2010; accepted 20 September 2010.

(Associate Editor: Robin Taylor)

INTRODUCTION

There are different terms around the world for bushfires, which are also called wildfires, forest fires, vegetation fires or brushfires. In this paper, we will use the term bushfire (mainly used in Australia), by which we mean fire that occurs in forest, scrub or grassland anywhere in the world.

Bushfires occur on every continent except Antarctica. Bushfire smoke has always been a part of Australia and Asia since human settlement. Fires mainly occur in climates where there are long dry spells. The Asia-Pacific region has experienced some of the worst bushfire events on record. For example, the 1997 Indonesian fires burnt over 5 million hectares, and 173 people lost their lives in the 'Black Saturday' 2009 fires in Australia. Health effects of bushfires go beyond the immediate severe health effects, such as loss of life or burns. During bushfires, large areas of land can be covered in layers of smoke, hundreds of kilometres away from the actual fires (Fig. 1), potentially involving major cities and exposing millions of people to bushfire smoke. Even small increases in risk could cause large public health problems.

Bushfire smoke, like background urban air pollution, consists of pollutants which can affect health. It contains large amounts of particulate matter (PM), but also polycyclic aromatic hydrocarbons, carbon monoxide, aldehydes, organic acids, volatile organic compounds and ozone following reactions in sunlight.¹ The pollutant most consistently elevated due to bushfire smoke is PM,^{2,3} most commonly measured as PM₁₀ (particles with a median aerodynamic diameter less than 10 µm).

In the past decade, interest has shifted towards the smaller fraction of PM₁₀, that is, PM_{2.5} (particles with a median aerodynamic diameter less than 2.5 µm) which are able to penetrate deeper into the alveolar region of the lung and can potentially be more harmful.⁴ Currently, there is only an Advisory Reporting Standard for PM_{2.5} in Australia. However, very few studies on bushfire smoke and health have been able to directly assess effects of PM_{2.5}, simply because this fraction is not as easily measured as PM₁₀. Although 24-h gravimetric measurements of PM_{2.5} are feasible,




Figure 1 Satellite image made on 10/01/2003 showing south-east of Australia during a bushfire event. The locations of the fires are marked red (courtesy of Jeff Schmaltz, MODIS Rapid Response Team, National Aeronautics and Space Administration/Goddard Space Flight Center).




Figure 2 The air pollution health effects pyramid (adapted from American Thoracic Society 2000).⁴³

real time measurement requires more complex equipment such as Tapered Element Oscillating Microbalances.

Urban air pollution, and especially PM, has been associated with a wide range of health effects, including exacerbations of respiratory symptoms, declines in lung function, asthma attacks, emergency department (ED) presentations, hospital admissions and premature mortality.^{5,6} In fact, ambient PM air pollution has been estimated to be responsible for at least 0.8 million premature deaths and 6.4 million years of life lost worldwide.⁷ The health effects associated with particulate air pollution have been observed at levels well below current air quality standards. Table 1 presents the air quality standards for the countries covered in this review. During bushfire smoke epi-

sodes, PM concentrations several times above background urban concentrations can occur, with air quality standards commonly being exceeded. This then raises the question: If respiratory effects are observed well below current air quality standards, what health effects can be expected when air pollutant concentrations are significantly elevated due to bushfire smoke?

In this review, we will summarize the literature on the respiratory health effects of bushfire smoke. We will first summarize the literature available in the Asia-Pacific region, followed by studies from North America and Europe. Those studies with stronger study designs and rigorous analysis will be discussed in more detail.

We have only included peer-reviewed published studies which have investigated the association between bushfire events and respiratory morbidity. Some of these studies looked at other health outcomes such as cardiovascular effects, but for the purpose of this review we have only focused on respiratory outcomes. In Table 2, 3 and 4, we have included risk estimates for those studies which measured PM concentrations and related this to the health risks. The studies have used different methods to analyse their data ranging from simply comparing health events before and during a bushfire smoke episode, to more sophisticated statistical models. Case-crossover studies and time series represent different approaches to define associations between exposure to ambient air pollution and health events over time.²⁰

AUSTRALIA

Around Australia, studies of the effect of bushfire smoke on respiratory health have been done in

Table 1 Air quality standard for PM for countries where studies have been conducted and the WHO AQG

	PM ₁₀	Allowable exceptions	PM _{2.5}	Allowable exceptions
WHO AQG [†]	50 µg/µm ³ avg per day 20 µg/m ³ avg per year	—	25 µg/µm ³ avg per day 10 µg/m ³ avg per year	—
Australia	50 µg/m ³ avg per day	5 days per year	25 µg/m ³ avg per day [‡] 8 µg/m ³ avg per year [‡]	—
Indonesia	150 µg/m ³ avg per day	—	—	—
Malaysia	150 µg/m ³ avg per day 50 µg/m ³ avg per year	—	—	—
Singapore	150 µg/m ³ avg per day	—	35 µg/m ³ avg per day 15 µg/m ³ avg per year	—
European Union [§]	50 µg/m ³ avg per day 40 µg/m ³ avg per year	35 days per year	—	—
USA	150 µg/m ³ avg per day	1 day per year (on average over 3 years)	35 µg/m ³ avg per day 15 µg/m ³ avg per year	The 3 year average of the 98th percentile of 24-h PM is not allowed to exceed 35 µg/m ³

[†] The World Health Organisation assists countries in reducing the health effects of air pollution by providing guidance to countries in setting their standards.

[‡] PM_{2.5} is only an advisory reporting standard in Australia.

[§] Lithuania has adopted the European Union limit value.

Avg, average; PM, particulate matter; PM_{2.5}, particles with a median aerodynamic diameter less than 2.5 µm; PM₁₀, particles with a median aerodynamic diameter less than 10 µm; WHO AQG, World Health Organisation Air Quality Guidelines.⁸

Melbourne,⁹ Sydney,^{10–14} Brisbane²¹ and Darwin (Table 2).^{16–19}

Victoria

As for most of the Australian studies, Tham *et al.*⁹ used a time series approach to investigate the effects of PM₁₀, Airborne Particle Index (API representing fine particulate pollution 0.1–1 µm in diameter) and ozone on respiratory-related ED visits and hospital admissions during a 6-month bushfire season (2002–2003). Figure 1 shows a satellite image of the extensive smoke plume covering part of south-east Australia during this bushfire season. No associations were found in the Gippsland region of Victoria. However, the lack of association might be due to lack of statistical power as a result of a small population size or because those in rural areas lived further away from hospitals and were therefore more likely to go to primary health-care providers.

Nonetheless, this study found an association between PM₁₀ and ED visits after adjusting for day-of-the-week, temperature and humidity. An increase of 9.1 µg/m³ in PM₁₀ was associated with an increase in respiratory ED attendances of 1.8%. The associations with API and ozone were not significant. This study was done in a bushfire season period with fires burning for 2 of the 6-month study period. Although it is likely that a large amount of PM₁₀ was due to bushfire smoke, a limitation of this study was that the association found could not be attributed solely to bushfire PM₁₀. Two other longitudinal Australian

studies from Sydney¹⁴ and Brisbane²¹ covered several years and did try to attempt to separate bushfire PM₁₀ from background PM₁₀.

New South Wales

Several studies investigating the respiratory effects of bushfire smoke have been conducted in Sydney, with conflicting results. Two early studies investigated exposure to bushfire smoke in Sydney for the 1994 fires. They did not find an increase in emergency visits for asthma.^{10,11} The average daily air particulate concentration was not given, but daily PM₁₀ levels exceeded 150 µg/m³. The maximum hourly PM₁₀ reading in Sydney was 250 µg/m³.¹¹ Cooper and colleagues¹⁰ compared numbers of asthma ED visits in three inner-city hospitals for the month of January and divided this into three time periods. They did not find any significant difference between before, during or after the bushfires. Statistical power was an issue as the total number of ED presentations for the study period was less than 100. Complex statistical modelling was not used for this comparison and no confounding factors were considered.

The same bushfire period was investigated by Smith *et al.*¹¹ using ED presentations in western Sydney. The authors used two different analytical methods which would have been able to detect an increase in asthma presentations of 50% for the bushfire period. However, they also did not find any increase in ED presentations for the bushfire period for the week of the bushfires compared to the same

Table 2 Peer-reviewed studies investigating the association between bushfire smoke events and respiratory health effects in Australia

		Australia					
Location	Study	Study period	Study area	Exposure variable	Health outcome and study population	Analytical methodology	Study result
Victorian fires, 2002–2003	Tham <i>et al.</i> , 2009 ⁹	October 2002–April 2003	Melbourne and Gippsland region	PM ₁₀ , API	ED presentations and hospital admissions for respiratory disease	Time series analysis adjusting for temperature, humidity and day of the week	A 9.1 µg/m ³ increase in PM ₁₀ was associated with a 1.8% (95% CI: 0.4–3.3%) increase in respiratory-related ED presentations in Melbourne. No association with hospital admission was found after adjustment for confounders.
New South Wales fire, 1994	Cooper <i>et al.</i> , 1994 ¹⁰	January 1994	Sydney	—	ED presentations for asthma at three inner-city hospitals	No details given	No increase in asthma presentations compared with before the event.
	Smith <i>et al.</i> , 1996 ¹¹	January 1994	Western Sydney	PM ₁₀	ED presentations for asthma at seven hospitals	Comparisons between case and control periods and time series analysis ¹	No association between asthma presentations and PM ₁₀ from bushfire smoke.
	Jalaludin <i>et al.</i> , 2000 ¹²	January 1994	Sydney	PM ₁₀	PEFR in 32 children with wheeze	Time series analysis adjusted for temperature, humidity, time spent outdoors and pollen count	No association between the bushfire period or daily PM ₁₀ and evening PEFR was found.
	Jalaludin <i>et al.</i> , 2004 ¹³	January 1994	Sydney	PM ₁₀	Evening recorded symptoms (wheeze, dry cough and wet cough) and beta agonist use in 32 children from three schools in Sydney with a history of wheeze	Time series analysis adjusted for temperature, humidity, number of hours spend outdoors, pollen count and inhaled corticosteroid use	Increase of 35 µg/m ³ same day bushfire PM ₁₀ was associated with an evening wet cough (OR 2.04, 95% CI: 1.37–3.04). None of the other symptoms were significant and neither was medication use.
New South Wales fires 1994–2002	Morgan <i>et al.</i> , 2010 ¹⁴	1994–2002	Sydney	PM ₁₀	Hospital admissions for respiratory disease	Time series analysis adjusting for temperature, humidity, day of week and flu epidemic	A 10 µg/m ³ increase in bushfire PM ₁₀ was associated with a 1.24% (95% CI: 0.22–2.27%) increase in all respiratory hospital admissions.
Queensland fires	Chen <i>et al.</i> , 2006 ¹⁵	July 1997–December 2000	Brisbane	PM ₁₀	Hospital admissions for respiratory disease	Time series analysis adjusted for average temperature, day of the week, seasonality, long-term trends (years) and influenza	Comparing hospital admission on high PM ₁₀ days (>20 µg/m ³) to low PM ₁₀ days (<15 µg/m ³) showed an increase in respiratory hospital admissions of 19% (95% CI: 9–30%) for bushfire days and of 13% (95% CI: 6–23%) for background days.

Table 2 Continued

Australia							
Location	Study	Study period	Study area	Exposure variable	Health outcome and study population	Analytical methodology	Study result
Northern Territory fires	Johnston <i>et al.</i> , 2002 ¹⁶	April–October 2000	Darwin	PM ₁₀	ED presentations for asthma	Time series analysis adjusting for weekly influenza and weekday versus weekend day	There was an increase in asthma presentations when average daily PM ₁₀ ≥ 40 µg/m ³ was compared to the baseline category of <10 µg/m ³ (Rate ratio: 2.39, 95% CI: 1.46–3.90). An increase of 10 µg/m ³ in PM ₁₀ was non-significantly associated with hospital admissions for all respiratory conditions (OR 1.08, 95% CI: 0.98–1.18), and significantly associated with COPD admissions (OR 1.21, 95% CI: 1.00–1.47). For indigenous people all effect sizes were larger, in particular for COPD presentations (OR 1.98, 95% CI: 1.10–3.59).
	Johnston <i>et al.</i> , 2007 ¹⁷	April–November of 2000, 2004 and 2005	Darwin	PM ₁₀	Hospital admissions for respiratory diseases	Case-crossover analysis adjusted for weekly influenza rate, days with rainfall >5 mm, temperature and humidity for same day and previous 3 days and public holidays	
	Hanigan <i>et al.</i> , 2008 ¹⁸	April–November from 1996 to 2005	Darwin	PM ₁₀ (measured and estimated)	Hospital admissions for respiratory diseases	Time series analysis adjusted for indigenous status, influenza epidemics and holidays	An increase of 10 µg/m ³ same day (estimated) PM ₁₀ was associated with a non-significant 4.81% (95% CI: –1.04–11.01%) increase in total respiratory admissions. However this association was significant for indigenous people with a 9.40% increase (95% CI: 1.04–18.46%).
	Johnston <i>et al.</i> , 2006 ²³	April–October 2004	Darwin	PM ₁₀ and PM _{2.5}	Asthma symptoms and medication use in 251 adults and children with asthma	Time series analysis adjusted for temperature, humidity, pollen and spore counts, weekly influenza rates, weekends and holiday periods	An increase in 10 µg/m ³ was associated with an increase in risk of becoming symptomatic of 24% (95% CI: 11–39%) and 15% (95% CI: 7–23%) for PM ₁₀ and PM _{2.5} respectively. PM ₁₀ and PM _{2.5} were significantly associated with starting a course of oral steroids and PM _{2.5} with starting reliever medication.

[†] No confounding factors, like temperature and humidity included in the analysis. API, Airborne Particle Index; ED, emergency department; PEFR, peak expiratory flow rate; PM_{2.5}, particles with a median aerodynamic diameter less than 2.5 µm; PM₁₀, particles with a median aerodynamic diameter less than 10 µm.

Table 3 Peer-reviewed studies investigating the association between bushfire smoke events and respiratory health effects in Asia

Asia							
Location	Study	Study period	Study area	Exposure variable	Health outcome and study population	Analytical methodology	Study result
1997 Indonesian fires	Aditama, 2000 ²⁶	September 1997–June 1998	Indonesia	—	Routine data from governmental health facilities	Comparing cases between September 1997 and June 1998 with the same period in 1995–1996 ^{††}	Increase in cases of acute respiratory infection and bronchial asthma.
	Emmanuel, 2000 ²⁷	August–November 1997	Singapore	PM ₁₀	ED presentations, out-patient attendance, hospital admissions for haze-related conditions and mortality	Time series analysis adjusting for temperature, relative humidity, rainfall and wind speed	An increase in 100 µg/m ³ PM ₁₀ was significantly associated with a 12% increase in outpatient attendance for upper-respiratory tract illness, 19% for asthma and 26% for rhinitis. There were also increases in ED attendances for haze related conditions (no risks presented in paper). No increase in hospital admission or mortality was found.
	Mott <i>et al.</i> , 2005 ²⁸	January 1995–December 1998	Kuching, Malaysia	—	Hospital admission to seven hospitals for respiratory disease	Time series analysis comparing 1997 bushfire period with forecasted estimates using pre-fire periods, adjusting for monthly seasonal components	Increase in respiratory hospitalizations during the bushfire period, particularly due to asthma.
	Sastry <i>et al.</i> , 2002 ²⁹	1996–1997	Kuala Lumpur and Kuching, Malaysia	PM ₁₀ [‡]	Mortality	Time series analysis [†]	Daily increase in haze levels associated with increased mortality rates for older individuals, although not significant for respiratory mortality. [‡]
	Kunii <i>et al.</i> , 2002 ²⁴	September 1997	Jambi, Indonesia	—	Respiratory symptoms in 492 children and adults	Reporting of percentage of respondents with respiratory symptoms ^{††}	A total of 91% of respondents reported respiratory problems due to the smoke haze caused by the bushfires.
	Brauer and Jamal, 1998 ³⁰	August and September 1997	Kuala Lumpur, Malaysia	—	Hospital admissions to major hospitals for asthma and respiratory infection	Comparing September admissions to August ^{††}	Increase in hospital admission for asthma and acute respiratory infection.
	Awang <i>et al.</i> , 2000 ³¹	September 1997	Selangor, Malaysia	—	Hospital admissions for respiratory disease	Comparing September figures to June figures [†]	Increase in hospital admissions for total respiratory disease, and in particular asthma and acute respiratory infection.
1994 Indonesian fires	Chew <i>et al.</i> , 1995 ³²	September–October 1995	Singapore	PM ₁₀	ED presentations for acute childhood asthma in children <12 years in two large hospitals in Singapore	Time series analysis adjusted for meteorological variables and 'other factors' (not further specified)	Increase in daily PM ₁₀ associated with increase in asthma emergency presentations (no risks presented in paper).

[†] No confounding factors, like temperature and humidity included in the analysis.

^{††} No statistical significance testing.

[‡] PM₁₀ was only used to identify high air pollution days (hence no risks presented for increases in PM).

ED, emergency department; PM, particulate matter; PM_{2.5}, particles with a median aerodynamic diameter less than 2.5 µm; PM₁₀, particles with a median aerodynamic diameter less than 10 µm.

Table 4 Peer-reviewed studies investigating the association between bushfire smoke events and respiratory health effects in North America and Europe

North America							
Location	Study	Study period	Study area	Exposure variable	Health outcome and study population	Analytical methodology	Study result
1987 Californian fires	Duclos <i>et al.</i> , 1990 ³³	August 1987	California	TSP, PM ₁₀	ED presentations to 15 hospitals in 6 Californian counties	Compared the 2.5 weeks of bushfire smoke to a control period [†]	Increase in ED presentations during the fire period for asthma, COPD, laryngitis, sinusitis and other respiratory infections.
1999 Californian fires	Mott <i>et al.</i> , 2002 ³⁴	August–November 1999	Humboldt County, California	—	Medical visits for respiratory illnesses to a local medical centre	Comparing the bushfire period visits to the previous year [†]	Medical visits for respiratory illnesses increased by 52% (217) visits during the fire period compared to the previous year.
2003 Californian fires	Viswanathan <i>et al.</i> , 2006 ³⁵	October–November 2003	San Diego County	PM ₁₀	ED presentations to 15 hospitals in San Diego County for respiratory diseases	Comparing 2 weeks during and following the bushfire to 1 week before the fire ^{††}	Increase in ED presentations during the fire period for asthma, respiratory problems (without fever) and smoke inhalation.
1998 Florida fires	Kunzli <i>et al.</i> , 2006 ³⁸	October–November 2003	California	PM ₁₀ and retrospective reporting on smell of fire indoors	Respiratory symptoms (including wheeze, cough and asthma attacks), medication usage and physician visits during the 2-week fire period for 4609 participants in the Children's Health Study in 16 communities	Mixed effects models adjusted for gender, ethnicity, educational levels of the parents, asthma status	Increase in all respiratory symptoms for those who reported an indoor fire smell of more than 6 days.
1998 Florida fires	Sorenson <i>et al.</i> , 1999 ^{39,40}	June–July 1998	Volusia and Flagler counties in Florida	—	ED presentations and hospital admissions at eight hospitals for respiratory conditions	Comparing bushfire period (1 June–6 July 1998) to the same period the year before ^{††}	ED presentations during the bushfire period increased by 91% for asthma (from 77 to 147), 132% for bronchitis with acute exacerbation (from 28 to 65), 37% for chest pain (from 218 to 299). Changes in the number of hospital admissions were minimal.
2003 British Columbia fires	Moore <i>et al.</i> , 2006 ⁴¹	1993–2003	Regions of Kelowna and Kamloops in British Columbia, Canada	PM ₁₀ and PM _{2.5}	Physician visits billed to respiratory disease diagnosis code	Comparing three weeks forest fire period with aggregated rates of same weeks in 10 previous years [†]	Significant increase in weekly physician visits for respiratory diseases during the fire period compared to same weeks in previous year for the Kelowna region. No significant effects were found for Kamloops (Kamloops experienced lower PM concentrations than Kelowna).
Europe							
2002 Lithuanian fires	Ovadnevaite <i>et al.</i> , 2006 ⁴²	August–September 2002	Vilnius City, Lithuania	—	Presentations at eight Vilnius health centres for respiratory diseases	Comparing increase in presentations from 1–18 September 2002 to July 2002 ^{††}	The number of presentations for respiratory diseases on average over all eight health centres was 3.1 times higher in September compared to July (ranging from 1.5 times in one health centre to 20.5 times in another).

[†] No confounding factors, like temperature and humidity included in the analysis.

^{††} No statistical significance testing.

ED, emergency department; PM, particulate matter; PM_{2.5}, particles with a median aerodynamic diameter less than 2.5 µm; PM₁₀, particles with a median aerodynamic diameter less than 10 µm. TSP, total suspended particles.

week the previous year. Considering the small effects found in other studies, there was not enough power to detect an effect.

Another study conducted during the same period investigated the effect of bushfire smoke on respiratory symptoms and peak expiratory flow rate (PEFR) in children.^{12,13} This longitudinal study was conducted in Sydney during January 1994 when children (average age 9 years) were measuring their PEFR and reporting on symptoms (wet cough, dry cough and wheezing) and medication use. Bushfire PM₁₀ was calculated by subtracting the average PM₁₀ concentration for January 1994 excluding the bushfire period from total daily PM₁₀. The PEFR analyses in 32 children with a history of wheeze found no association. An increase of 35 µg/m³ in same day bushfire PM₁₀ was significantly associated with a twofold increase in evening wet cough. Although no associations were found with dry cough, wheezing or medication use, this might have been expected as these symptoms have shown to be associated with urban background PM₁₀. The study findings may have been influenced by limited statistical power.

A more recently published study by Morgan *et al.*¹⁴ investigated respiratory hospital admissions with an extended time series analysis from 1994 to 2002. They divided PM₁₀ into bushfire PM₁₀ and 'non-bushfire' (background) PM₁₀. It was assumed that on days where city-wide 24-h average PM₁₀ was greater than the 99th percentile for the study period, the PM₁₀ was primarily from bushfires. On these days bushfire PM₁₀ was calculated to be the difference between total PM₁₀ and estimated background PM₁₀. Over the 8.5-year study period, 32 bushfire days were identified, resulting from 14 different bushfire events. Analyses were done for three age groups, and models adjusted for temperature, humidity, day of the week and presence of an influenza epidemic.

Both bushfire and background PM₁₀ were associated with small but significant increases in respiratory hospital admissions for Lag 0 (24-h average concentration on the day of the hospital admissions). However, no significant associations were found with 24-h average concentration on 1, 2 and 3 days before the hospital admission. When the analysis was restricted to respiratory admissions of people over 65 years of age, bushfire PM₁₀ showed a more consistent association than background PM₁₀. The greatest differences in effect between bushfire PM₁₀ and background PM₁₀ were in admissions for COPD among those over 65 years of age and for asthma admissions for those between 15 and 64 years of age. No associations were found with background PM₁₀, but significant associations were found for both outcomes with bushfire PM₁₀. The COPD admissions for those over 65 years were consistently associated with bushfire PM₁₀, with a 10 µg/m³ increase in bushfire smoke resulting in an increase in hospital admissions of 3.3%. Asthma admissions for the 15–64 year age group increased by 5.0% for a 10 µg/m³ increase in bushfire PM₁₀. Perhaps surprisingly, no association was found for asthma admissions among children 1–14 years of age.

In the same study, no association was found between bushfire PM₁₀ and respiratory mortality.

However, the number of daily respiratory deaths in Sydney was low (with a median of 5) and this limited the power of the analysis to find an effect even if one was present. Background PM₁₀ did not show an association with respiratory mortality either, even though this analysis had more power to detect an effect.

Queensland

Another time series study from Brisbane investigated the association between bushfire PM₁₀ on respiratory hospital admissions.²¹ The authors used three categories for average daily PM₁₀ in the analysis. A bushfire day included an event where more than 1 ha of area was burnt in the Brisbane region. During the study period there were 452 bushfire days (35% of all days). The daily average PM₁₀ concentration over 452 bushfire days was 18 µg/m³ (range 8–61 µg/m³) and for the 828 background days 15 µg/m³ (range 5–58 µg/m³). These mean values were well below the current Australian air quality standards (see Table 1). There was an increase in respiratory hospital admissions of 19% for bushfire days and of 13% for days without the presence of bushfire smoke. A limitation of this study is that data were obtained from a PM₁₀ monitor which was upwind of most bushfires, therefore probably underestimated the exposure on bushfire days. Nonetheless, on bushfire days the association with respiratory hospital admission was at least as great as with urban background PM₁₀. There was a suggestion of a greater effect of bushfire PM₁₀, but this would need further confirmation.

Northern Territory

Darwin provides a unique opportunity to study the effects of bushfire PM because approximately 95% of PM₁₀ is due to fire smoke for up to 8 months a year, and only 5% is made up of industrial and motor vehicle emissions.²² The first study examined the association between ED presentations for asthma and PM₁₀ during the 2000 bushfire period.¹⁶ The average daily PM₁₀ was 21 µg/m³ (range 2–70 µg/m³), and there were 7 days in which the National Environment Protection Measures (NEPM) for PM₁₀ was exceeded during this period. Four categories of PM₁₀ concentrations were studied and analyses were adjusted for weekly rates of influenza and for weekday versus weekend. There was a 1.2-fold increase in asthma presentations for every 10 µg/m³ increase in PM₁₀.

Another study by Johnston and colleagues¹⁷ from Darwin investigated the association between hospital admissions for all respiratory diseases and bushfires during the 2000, 2004 and 2005 seasons combined. Unfortunately, no PM₁₀ measurement data were available for the years 2001 to 2003. This is the only study on bushfire smoke which has used the case-crossover design that eliminates confounding by day of week and seasonal trends in the exposure variables. An increase of 10 µg/m³ in PM₁₀ was most strongly associated with COPD and asthma. For indigenous people all effect sizes were larger, in particular for COPD presentations.

Hospital admission for respiratory diseases in Darwin were further investigated by Hanigan *et al.*¹⁸ who used indirect estimates of PM₁₀, derived from visibility data for the bushfire periods in Darwin (when no direct PM measurements were available). They analysed 1996 to 2005 data and found that a 10 µg/m³ increase in estimated PM₁₀ was associated with a non-significant increase in total respiratory admissions on the same day. However, this association was significant for indigenous people with a 9.4% increase. Asthma, COPD and respiratory infections did not show significant associations, but this could be due to small numbers and a lack of statistical power.

A cohort study from Darwin followed 251 adults and children with asthma over the 2004 bushfire period.²³ Participants kept a daily diary where they recorded their asthma symptoms, medication use, missed school or work days and health-care visits for asthma. Analyses were adjusted for temperature, humidity, pollen and spore counts, weekly influenza rates, weekends and holiday periods. PM₁₀ levels were rather low and there were no occasions when the NEPM were exceeded during the study period. Significant associations were found with PM₁₀ and PM_{2.5} and development of symptoms. An increase in 10 µg/m³ was associated with an increase in asthma symptoms of 24% for PM₁₀ and 15% for PM_{2.5}. PM₁₀ and PM_{2.5} were also significantly associated with the need for a course of oral steroids and PM_{2.5} with increased reliever medication. Due to the low PM concentrations, high correlations and small effects it was not possible to distinguish between PM₁₀ and PM_{2.5} effects.

ASIA

One of the largest forest fire episodes for which effects on community health have been reported occurred in Indonesia in 1997. Approximately 5.3 million hectares were burned during these fires²⁴ and the severity and extent of the associated smoke haze affected about 300 million people across the region.²⁵ Several studies investigated the effect on respiratory health of the populations living in cities across South-East Asia (Table 3). However, only one study was done in Indonesia, where the actual fires occurred.²⁶ Routine data from government health facilities were used to compare cases between September 1997 and June 1998 with the same period in 1995–1996 and increases were reported for acute respiratory infection and bronchial asthma. For example, in south Kalimantan the number of acute respiratory infections increased by 1.8 times, and in south Sumatra this was 3.8 times.

A further seven published studies were done outside Indonesia.^{24,26–32} For example, across the strait in Singapore, an increase of 100 µg/m³ in PM₁₀ was associated with a 19% increase in asthma and 12% increase in upper respiratory tract illnesses. However, this study did not find an association with hospital admissions or mortality.²⁷ A study looking at hospital admission in the Kuching region of Malaysia found significant increases in respiratory hospitalizations particularly due to asthma, during the bushfire period

compared to previous year²⁸ but no air pollution concentrations were measured in this study.

As Table 3 shows, even studies done outside Indonesia and some distance away from the fires found an effect on respiratory health during smoke haze episodes in South-East Asia. Increases were found for emergency admissions for all respiratory diseases or asthma alone, and in most cities increases were found for hospital admissions for respiratory diseases but there was no significant effect on respiratory mortality.

NORTH AMERICA

Hospital emergency or local medical centre visits increased after the California wildfires in 1987,³³ 1999³⁴ and 2003.³⁵ There was a unique opportunity during the 2003 California fires, when regions that participated in the Children's Health Study^{36,37} were affected by bushfire smoke. The resulting analysis found that the smell of bushfire penetrated indoors for more than 6 days, respiratory symptoms (sore throat, cough, bronchitis, wheezing and asthma attacks) and medication usage were significantly increased. An important limitation was that both exposure and outcome were reported by participants which could lead to interrelated reporting. Associations were strongest for those without asthma, perhaps because those with asthma were more likely to stay indoors and wear masks during the smoke episode. This study measured 5-day average PM₁₀ concentrations during the fire period in each of the 16 participating communities, with concentrations ranging from 30 µg/m³ to 210 µg/m³. An increase in 5-day average PM₁₀ was associated with dry cough, sore throat and medication use, but no significant associations were found with wet cough, bronchitis, wheezing or asthma attacks.³⁸

During the 1998 fires in Florida, increases in weekly respiratory ED presentations were found compared to the same weeks in 1997. This effect was strongest for asthma and acute exacerbations of bronchitis. However, this was not associated with increases in weekly hospital admissions, although the number of hospital admissions during the bushfire period was small. The results from this study should be interpreted with caution as no air quality measurements were reported.^{39,40}

The 2003 fires in British Columbia resulted in an increase in weekly rates of physician visits for respiratory diseases in the Kelowna region, but not in the Kamloops region.⁴¹ The absence of effects in Kamloops, could be due to less power to detect an effect as the population is smaller than in Kelowna. Alternatively, it could be due to lower levels of PM₁₀ in Kamloops (maximum daily average 150 µg/m³ cf. 200 µg/m³). In Kelowna, the largest increases in physician visits for respiratory diseases were observed in the fifth week after the fires began, when PM₁₀ levels returned to background levels. This could be either because the respiratory symptoms may take several weeks to develop, or suggest that patients may initially try to manage their symptoms on their own before visiting a physician.

EUROPE

In a study from Lithuania where there was a bushfire episode in 2001, the number of visits to eight health centres in Vilnius for respiratory diseases increased 2.8-fold compared to 1 month when there were no fires.⁴² Given the serious bushfires in Greece and elsewhere in southern Europe in recent years, it is surprising that more research has not been reported from this part of the world (Table 4).

DISCUSSION

The peer-reviewed literature to date suggests that there is a modest association between bushfire smoke and respiratory health. For example, a comprehensive Australian study over several years found an increase in hospital admissions of 1.2% for each 10 µg/m³ increase in PM₁₀ from bushfire smoke.¹⁴ Estimating the overall effects of bushfire smoke on asthma, COPD or mortality is complicated because of the different methods used in the studies.

The association between exposure to bushfire smoke and respiratory morbidity is consistent with the associations found with urban air pollution. All studies looking at ED presentations found associations, with either increases during the bushfire period compared to a control period, or when appropriate statistical modelling was used and confounding factors were taken into account. However, only a few studies have attempted to differentiate between the effects of bushfire PM₁₀ and background PM₁₀.^{14,21} It would be too difficult to perform a meta-analysis of the studies reviewed here, due to the large differences in study design and statistical methods used. Most studies did not directly measure PM concentrations or used estimates of PM. Some studies investigated all respiratory effects and others only looked at asthma. The time period of analysis also differed between studies. Of all the studies carried out, only three from Sydney did not find an association.^{10–12} All these have statistical limitations. However we accept the likelihood of publication bias, with positive studies being more likely to be published.

A limitation of studies investigating the health effects of bushfire smoke is that the smoke episodes are usually of short duration. Statistical power could be increased by studying larger populations and/or over longer periods. For example, Tham *et al.*⁹ investigated respiratory hospital admissions both in Melbourne and Gippsland during a bushfire season. The average PM₁₀ concentrations over the study period for the two locations were similar, but the maximum daily concentrations in Gippsland were larger. The study found a small but significant effect of PM₁₀ in Melbourne, but not in Gippsland, likely due to the much larger population in Melbourne and hence greater power to detect an effect. A similar result was seen in two regions in Canada.⁴¹ A significant increase in respiratory physician visits during the fire period was found in the town with the larger population and

higher PM₁₀ concentrations. Several studies with rather low PM₁₀ concentrations did find an effect, but these studies were carried out over years rather than weeks or months.^{14,16,21}

The studies of respiratory effects of bushfires have usually focused on outcomes that can be obtained shortly after the bushfire smoke event, simply because episodes are difficult to predict in advance. As a result most research has focused on emergency presentations, hospital admissions or mortality. However, these outcomes are at the tip of the health effects pyramid (Fig. 2). Only a few studies have been carried out on more subtle outcomes, often in relation to studies which were ongoing at the time of an event.^{12,13,38} Another issue with studying health effects of bushfire smoke is that they tend to occur in rural areas and if there is not a large city nearby, the population is usually not large enough to determine with confidence whether an effect has occurred.

Some ambient air pollutants are regulated because they are thought to be harmful to health at certain concentrations. In Australia, the National Environment Protection Council sets ambient air quality goals for six pollutants in the NEPM for ambient air quality. These six pollutants are ozone, particles, carbon monoxide, nitrogen dioxide, sulphur dioxide and lead. In Australia, the NEPM for acceptable daily average PM₁₀ is 50 µg/m³. Currently, the NEPM PM₁₀ standard is allowed to be exceeded 5 days per year, to accommodate for bushfire smoke. However, during bushfire episodes the standards are often exceeded more than the 'allowable' 5 times.³ For example, the PM₁₀ NEPM target was exceeded on a total of 6 days in the 7-month bushfire period in Darwin in 2000.¹⁶

There are very clear effects on mortality in urban air pollution studies,⁵ but these are not found in the bushfire smoke studies. This could very well be due to a lack of power in mortality analyses. The number of deaths are smaller than the number of ED presentations or hospital admissions (Fig. 2). In addition, bushfire smoke episodes are usually of short duration, compared to the urban air pollution studies which are usually carried out over several years. For example, Vedal and Dutton⁴⁴ did not find an association with mortality and exposure to PM_{2.5} or PM₁₀. Even though they studied a population of 2 million people, very high PM concentrations due to wildfire smoke only occurred for 2 days. To be able to detect an effect in this study wildfire smoke PM_{2.5} would have to be more than 10 times more toxic than urban background PM_{2.5}.⁴⁵ Hanninen *et al.*⁴⁶ found a consistent, but non-significant association between mortality and PM_{2.5} during a 2-week bushfire smoke episode. This suggested that an effect could be present, but the power of the study was such that significance could not be achieved.

From the evidence available to date, it is difficult to determine whether bushfire smoke PM₁₀ is more toxic than background PM₁₀. However, the two studies that attempted to separate bushfire PM₁₀ from background PM₁₀ suggest that bushfire PM₁₀ may have a greater impact on respiratory health than urban background PM₁₀,^{14,21} in particular for hospital admissions

Box 1 Bushfire: asthma management tips, adapted from the Asthma Foundation of Victoria Information Sheet.

- If you live in a fire danger area, or are likely to be visiting one, discuss the dangers with your local general practitioner and update your personal written Asthma Action Plan accordingly. You should also include asthma management in your fire safety survival plan.
- Always follow your personal written Asthma Action Plan. These instructions help you, your family, carers and neighbours recognize signs of worsening asthma and what action to take to prevent further deterioration.
- Always ensure you have plenty of medication on hand, particularly blue reliever medication. Continue use of your preventer as well.
- If you decide to evacuate, make sure your blue reliever medication goes with you.
- Listen to your local radio station or watch television for updates on fire and smoke conditions in the area.
- When smoke is in the air but a fire is not directly threatening you, minimize the time you spend outdoors in the smoke. Where practical, stay indoors and close all windows and doors.
- Turn on your air conditioner (if you have one) and switch it to 'recycle' or 'recirculate'. This will reduce the amount of smoke coming inside.
- If possible stay in air-conditioned premises. If your home gets uncomfortably hot or is letting in outside air, take an air-conditioned break at a local community library or shopping centre.
- If you are particularly susceptible to bushfire smoke, consider staying with a friend or relative whose house has clean indoor air.
- Everyone should rest indoors and avoid outdoor activity as much as possible. Increased physical activity causes deeper breathing, and a greater number of fine particles are breathed deep into the lungs.
- If the bushfire event lasts a long time, take advantage of any breaks in smoky conditions to air out your home, but remember to close off the house again when conditions deteriorate.
- If you develop symptoms such as shortness of breath, coughing or wheezing, chest tightness follow your written Asthma Action plan. If your reliever medication is not reducing your asthma symptoms, seek medical attention without delay and start Asthma First Aid.

in those over 65 years of age, and asthma admissions in younger adults.¹⁴ These results are consistent with a review of studies of PM₁₀ and asthma that found greater risks when wood combustion was considered to be the main source of PM.⁴⁷

Bushfires are expected to increase in the future as a result of climate change, and this will result in a larger public health issue. Only a few thorough population studies have been carried out on the issue of bushfire smoke and respiratory health, and more research is needed to be able to properly advise policy makers and clinicians of appropriate public health messages and measures to put in place. However, the evidence in this review supports the current advice given by the Asthma Foundation (Box 1).

REFERENCES

- 1 Malilay J. A Review of Factors Affecting the Human Health Impacts of Air Pollutants from Forest Fires. In: Gsoh KT, Schwela D, Goldammer JG (eds). *Health guidelines for vegetation fire events—background papers*. World Health Organisation, Geneva, 1999; 258–83.
- 2 Brauer M. Health impacts of biomass air pollution. In: Goh KT, Schwela D, Goldammer JG (eds). *Health guidelines for vegetation fire events—background papers*. World Health Organisation, Geneva, 1999; 186–254.
- 3 Reisen F, Brown SK. Implications for community health from exposure to bushfire air toxics. *Environ. Chem.* 2006; **3**: 235–43.
- 4 Englert N. Fine particles and human health—a review of epidemiological studies. *Toxicol. Lett.* 2004; **149**: 235–42.
- 5 Pope CA, 3rd, Dockery DW. Health effects of fine particulate air pollution: lines that connect. *J. Air Waste Manag. Assoc.* 2006; **56**: 709–42.
- 6 HEI Panel on the Health Effects of Traffic-Related Air Pollution. *Traffic-Related Air Pollution: A Critical Review of the Literature on Emissions, Exposure, and Health Effects*. HEI Special Report 17. Health Effects Institute, Boston, MA, 2010.
- 7 Cohen AJ, Ross Anderson H, Ostro B *et al.* The global burden of disease due to outdoor air pollution. *J. Toxicol. Environ. Health A* 2005; **68**: 1301–7.
- 8 World Health Organization. *WHO Air Quality Guidelines for Particulate Matter, Ozone, Nitrogen Dioxide and Sulfur Dioxide. Global Update 2005. Summary of Risk Assessment*. World Health Organization, Geneva, 2006.
- 9 Tham R, Erbas B, Akram M *et al.* The impact of smoke on respiratory hospital outcomes during the 2002–2003 bushfire season, Victoria, Australia. *Respirology* 2009; **14**: 69–75.
- 10 Cooper CW, Mira M, Danforth M *et al.* Acute exacerbations of asthma and bushfires. *Lancet* 1994; **343** (8911): 1509.
- 11 Smith MA, Jalaludin B, Byles JE *et al.* Asthma presentations to emergency departments in western Sydney during the January 1994 Bushfires. *Int. J. Epidemiol.* 1996; **25**: 1227–36.
- 12 Jalaludin B, Smith M, O'Toole B *et al.* Acute effects of bushfires on peak expiratory flow rates in children with wheeze: a time series analysis. *Aust. N. Z. J. Public Health* 2000; **24**: 174–7.
- 13 Jalaludin B, O'Toole B, Morgan G *et al.* Acute effects of bushfires on respiratory symptoms and medication use in children with wheeze in Sydney, Australia. *Environ. Health* 2004; **4**: 20–9.
- 14 Morgan G, Sheppard V, Khalaj B *et al.* Effects of bushfire smoke on daily mortality and hospital admissions in Sydney, Australia. *Epidemiology* 2010; **21**: 47–55.
- 15 Chen L, Mengersen K, Tong S. Spatiotemporal relationship between particle air pollution and respiratory emergency hospital admissions in Brisbane, Australia. *Sci Total Environ* 2007; **373**: 57–67.

- 16 Johnston FH, Kavanagh AM, Bowman DM *et al.* Exposure to bushfire smoke and asthma: an ecological study. *Med. J. Aust.* 2002; **176**(11): 535–8.
- 17 Johnston FH, Bailie RS, Pilotto LS *et al.* Ambient biomass smoke and cardio-respiratory hospital admissions in Darwin, Australia. *BMC Public Health* 2007; **7**: 240.
- 18 Hanigan IC, Johnston FH, Morgan GG. Vegetation fire smoke, indigenous status and cardio-respiratory hospital admissions in Darwin, Australia, 1996–2005: a time-series study. *Environ. Health* 2008; **7**: 42.
- 19 Johnston FH, Bailie R, Pilotto L. Bushfire smoke and respiratory hospital admissions in the Australian monsoon tropics. *Aust Epidemiol* 2006; **13**: 65.
- 20 Rothman KJ, Greenland S, Lash TL. *Modern Epidemiology*. Lippincott Williams & Wilkins, Philadelphia, PA, 2008.
- 21 Chen L, Verrall K, Tong S. Air particulate pollution due to bushfires and respiratory hospital admissions in Brisbane, Australia. *Int. J. Environ. Health Res.* 2006; **16**: 181–91.
- 22 Gras J, Ayres G, Keywood M *et al.* *A Pilot Study of Air Quality in Darwin, NT*. Northern Territory Government, Department of Lands Planning and Environment, CSIRO Atmospheric Research, Aspendale, 2001.
- 23 Johnston FH, Webby RJ, Pilotto LS *et al.* Vegetation fires, particulate air pollution and asthma: a panel study in the Australian monsoon tropics. *Int. J. Environ. Health Res.* 2006; **16**: 391–404.
- 24 Kunii O, Kanagawa S, Yajima I *et al.* The 1997 haze disaster in Indonesia: its air quality and health effects. *Arch. Environ. Health* 2002; **57**: 16–22.
- 25 Afroz R, Hassan MN, Ibrahim NA. Review of air pollution and health impacts in Malaysia. *Environ. Res.* 2003; **92**: 71–7.
- 26 Aditama TY. Impact of haze from forest fire to respiratory health: Indonesian experience. *Respirology* 2000; **5**: 169–74.
- 27 Emmanuel SC. Impact to lung health of haze from forest fires: the Singapore experience. *Respirology* 2000; **5**: 175–82.
- 28 Mott JA, Mannino DM, Alverson CJ *et al.* Cardiorespiratory hospitalizations associated with smoke exposure during the 1997, Southeast Asian forest fires. *Int. J. Hyg. Environ. Health* 2005; **208**: 75–85.
- 29 Sastry N. Forest fires, air pollution, and mortality in southeast Asia. *Demography* 2002; **39**: 1–23.
- 30 Brauer M, Hisham-Hashim J. Fires in Indonesia: crisis and reaction. *Environ. Sci. Technol.* 1998; **32** (17): 404A–7A.
- 31 Awang MB, Jaafar AB, Abdullah AM *et al.* Air quality in Malaysia: impacts, management issues and future challenges. *Respirology* 2000; **5**: 183–96.
- 32 Chew FT, Ooi BC, Hui JK *et al.* Singapore's haze and acute asthma in children. *Lancet* 1995; **346** (8987): 1427.
- 33 Duclos P, Sanderson LM, Lippsett M. The 1987 forest fire disaster in California: assessment of emergency room visits. *Arch. Environ. Health* 1990; **45**: 53–8.
- 34 Mott JA, Meyer P, Mannino D *et al.* Wildland forest fire smoke: health effects and intervention evaluation, Hoopa, California, 1999. *West. J. Med.* 2002; **176**: 157–62.
- 35 Viswanathan S, Eria L, Diunugala N *et al.* An analysis of effects of San Diego wildfire on ambient air quality. *J. Air Waste Manag. Assoc.* 2006; **56**: 56–67.
- 36 Peters JM, Avol E, Gauderman WJ *et al.* A study of twelve Southern California communities with differing levels and types of air pollution. II. Effects on pulmonary function. *Am. J. Respir. Crit. Care Med.* 1999; **159**: 768–75.
- 37 Peters JM, Avol E, Navidi W *et al.* A study of twelve Southern California communities with differing levels and types of air pollution. I. Prevalence of respiratory morbidity. *Am. J. Respir. Crit. Care Med.* 1999; **159**: 760–7.
- 38 Kunzli N, Avol E, Wu J *et al.* Health effects of the 2003 Southern California wildfires on children. *Am. J. Respir. Crit. Care Med.* 2006; **174**: 1221–8.
- 39 From the Centers for Disease Control and Prevention. Surveillance of morbidity during wildfires—Central Florida, 1998. *JAMA* 1999; **281**: 789–90.
- 40 Sorensen B, Fuss M, Mulla Z *et al.* Surveillance of morbidity during wildfires—Central Florida, 1998. *MMWR Morb. Mortal. Wkly. Rep.* 1999; **48**: 78–9.
- 41 Moore D, Copes R, Fisk R *et al.* Population health effects of air quality changes due to forest fires in British Columbia in 2003: estimates from physician-visit billing data. *Can. J. Public Health* 2006; **97**: 105–8.
- 42 Ovadnevaite J, Kvietkus K, Marsalka A. 2002 summer fires in Lithuania: impact on the Vilnius city air quality and the inhabitants health. *Sci. Total Environ.* 2006; **356**: 11–21.
- 43 American Thoracic Society. What constitutes an adverse health effect of air pollution? Official statement of the American Thoracic Society. *Am. J. Respir. Crit. Care Med.* 2000; **161** (2 Pt 1): 665–73.
- 44 Vedal S, Dutton SJ. Wildfire air pollution and daily mortality in a large urban area. *Environ. Res.* 2006; **102**: 29–35.
- 45 Hanninen O. Challenges in estimating the health effects of biomass smoke—response to Sverre Vedal and Steven J. Dutton: wildfire air pollution and daily mortality in a large urban area. *Environ. Res.* 2008; **106**: 423–4; discussion 425.
- 46 Hanninen OO, Salonen RO, Koistinen K *et al.* Population exposure to fine particles and estimated excess mortality in Finland from an East European wildfire episode. *J. Expo. Sci. Environ. Epidemiol.* 2009; **19**: 414–22.
- 47 Boman BC, Forsberg AB, Jarvholm BG. Adverse health effects from ambient air pollution in relation to residential wood combustion in modern society. *Scand. J. Work Environ. Health* 2003; **29**: 251–60.