

On COVID19 epidemiology, with application to wildland fire management practices

Karin Riley, March 29, 2020

Epidemiologists are working with emerging information in order to understand the spread of COVID-19, who is at high risk, the fatality rate, etc. The information below is the best available as of this writing, but is likely to change rapidly, making frequent updates necessary. It is too soon for much information to be peer-reviewed and published, though more is appearing daily. Much of the literature currently appears in the form of fact sheets, letters to journals, and blogs. For other items, reliance on conversations with epidemiologists and media reports is necessary.

As the situation and what is known about the disease is changing rapidly, frequent update of this white paper is recommended. Because little is known with certainty about transmission, fatality rates, etc., one author recommends: “It is possible to communicate uncertainty about the underlying science, apply the precautionary approach where uncertainty exists, and implement well-reasoned decisions about how best to limit the dissemination of COVID-19” (Brosseau 2020).


- **Disease emergence, spread, fatality rate, and mutation**
 - The disease emerged in late December 2019 in Wuhan, China (Tang et al In Press). The virus is called SARS-CoV-2 and it causes the disease COVID-19 or generically “the coronavirus”; however, the latter term is confusing as it encompasses a family of viruses including the common cold and SARS-CoV-1 that caused the previous SARS epidemic (Dr. Emily Ricotta, NIH, personal communication, 3/27/20).
 - As of Sunday, March 29, 685,623 confirmed cases of COVID-19 were reported globally, with at least 32,137 people dying of the disease, according to Johns Hopkins University (Johns Hopkins 2020). Another 145,706 people had recovered. The fatality rate in known cases then currently hovers around 18% globally, if we calculate fatality rate as $(\text{dead})/(\text{dead} + \text{recovered})$, a method recommended for emerging infectious diseases (Ghani et al 2005)
 - The situation is changing rapidly and the number of infected, recovered, and dead can be tracked online (e.g. <https://systems.jhu.edu/> or <https://www.worldometers.info/coronavirus/>)
 - The lack of availability of testing and the fact that many mild cases are not reported means that the current reach of the disease is not known, and the number affected are underestimated (e.g. Feng 2020). As many as 86% of cases may be undocumented (Li et al 2020). Therefore, the fatality rate given above is likely an overestimate, since more people are likely recovered but were not tested or reported. However, doctors report that patients are dying of COVID-19 without it being reported, so neither the number of dead or recovered can be known with any certainty (Prakash and Hall 2020).

- Testing on the Diamond Princess cruise ship indicated that almost half of cases were asymptomatic at the time of testing (46.5%) and 17.9% of infected people never developed symptoms (Moriarty et al 2020).
- Spread rate is currently exponential globally. Three months' time was required to reach 100,000 cases; the number doubled in 12 days (WHO 2020).
- The fatality rate varies with a number of factors, including access to medical care, whether high risk factors are present, and the patient's age (CDC 2020, personal communication with researchers working in the field).
- Severe symptoms and death can occur in any age group. Fatality rates in the US differ by age group and increase with age. People aged 85 years or older had a fatality rate of 10-27%, 3-11% among those 65-84 years of age, 1-3% in ages 55-64 years, <1% in those aged 20-54 years, and no fatalities recorded as of a March 18, 2020 report in those less than 19 years of age (CDC 2020).
- Over half of children in one Chinese study who contracted SARS-CoV-2 had radiologic findings consistent with pneumonia but no clinical presentation; one child in the study died (Lu et al 2020).
- COVID-19 likely poses the greatest threat to public health from a respiratory virus since the 1918 H1N1 influenza pandemic (Ferguson et al 2020).
- According to one group of scientists, after the SARS-CoV-2 virus that causes COVID-19 made the zoonotic jump from its animal host (likely a bat by way of a pangolin), the virus experienced rapid mutation (Tang et al In Press). Two distinct strains of the virus, known as the L and S strains, currently exist (Tang et al In Press). In some patients, mutations in the virus from both strains have been identified, meaning that co-infection or two subsequent infections are likely possible (Tang et al In Press). It is possible that a person infected with one strain would not be immune to the other. Therefore it may not be possible to consider a person who has had COVID-19 now immune to the disease, or if so, how long that immunity will last (Greenfieldboyce 2020).
- However, other scientists claim there is not evidence for two strains of the virus, that the virus is actually mutating slowly, and that one vaccine would likely be adequate to prevent infection (Achenbach 2020).
- Risk factors include age, gender (men are more likely to die of the disease), and co-morbidities, with the most common being hypertension, diabetes, and coronary heart disease (May 2020, Weiss and Murdoch 2020).
- The leading cause of death from COVID-19 is acute respiratory distress syndrome. An underrecognized second cause may be cytokine storm and hyperinflammation (Mehta et al 2020).
- **Disease transmission**
 - The SARS-CoV-2 virus is stable (meaning that it is still infectious) in aerosols for three hours, on cardboard up to 24 hours, and up to three days on stainless steel and plastic (van Doremalen et al 2020). The virus is still detectable even longer (for example, it is still detectable on cardboard up to 72 hours).

- The virus was found in cabins of symptomatic and asymptomatic people on the Diamond Princess cruise ship 17 days after the ship was vacated (Moriarty et al 2020).
- People who are infected with COVID-19 but have not been documented may be driving the spread of the disease, being responsible for 79% of documented cases in one study conducted during the early days of virus spread (Li et al 2020).
- Some health care professionals are voicing frustration with conflicting and sometimes shifting guidance from the Center for Disease Control (CDC) and World Health Organization (WHO) regarding how the disease transmits and how to protect oneself from contracting it (Brosseau 2020).
- According to the Chinese Center for Disease Control and Prevention, which likely has the greatest experience in dealing with cases of COVID-19 at this time, transmission of the disease is primarily through close contact and airborne droplets from coughing or sneezing, with transmission via aerosols possible in relatively closed environments when exposure to high concentrations is over some period of time (Chinese CDC 2020).
- The above suggests that the disease does not require droplets from coughing or sneezing in order to be transmitted, but may be transmitted through the air from breathing or talking or from touching contaminated objects (ScienceDaily 2020, Brosseau 2020).
- Transmission has been documented as far as 4.5 meters away and as long as 30 minutes after an infected person was present in a closed environment (Chen 2020).
- Prevention efforts should focus on droplet transmission rather than fomites (objects which may carry the disease), according to some researchers (Wölfel et al 2020).
- The more infectious particles one inhales, the greater the likelihood of contracting a disease (Brosseau 2020).
- Viability of the virus may be linked more strongly to the amount of time it has been in the air rather than the distance away from an infected person for viruses where aerosol or droplet transmission occurs (Brosseau 2020).
- Increased hand washing has been shown to be effective in reducing flu transmission in health care settings, but it may not be effectual in “community settings” based on a review of the available peer-reviewed literature (Moncion et al 2019). In other words, other measures besides hand washing may be needed to reduce disease transmission.
- Virus shedding may be at its highest level in the first five days as the disease and symptoms are developing, but it continues to be present in sputum through at least the first week during which symptoms are experienced (Wölfel et al 2020).
- Incubation period of the disease is variable, with the median being 5.1 days (Lauer et al 2020). A small number of cases (2.5%) took longer than 11.5 days for symptoms to appear.

- SARS-CoV-2 is present in urine and feces for a prolonged period after symptoms subside, with the virus appearing to replicate in the digestive tract (Wölfel et al 2020). It is not known whether the virus is contagious after travelling through the digestive tract (Wölfel et al 2020).
- Transmission of COVID-19 may be possible from direct contact with urine or feces, or particles aerosolized by urine or feces falling into a porta-potty or flushing of a toilet, however no cases have yet been documented to my knowledge (Chinese CDC 2020). Transmission through plumbing has not been ruled out and is suspected in a set of cases in Hong Kong (Regan 2020).
- High levels of COVID-19 can be found in rooms where workers remove protective clothing and in bathrooms, perhaps due to toilet flushing aerosolizing the virus (Brosseau 2020, Johnson et al 2013).
- Close contact is likely the primary risk factor for transmission, with close contact defined as being within 1 meter (Chinese CDC 2020).
- Travelling in a vehicle with others may increase likelihood of transmission, as the number of viruses may become concentrated in the air in this closed space.
- It is too soon to know how factors like heat, humidity, and pollution affect SARS-CoV-2, though research is currently underway.
 - However, research with previous SARS strains (CoV-P9) indicates that the virus remained infectious at temperatures between 39-100 degrees F for 2 hours or more, but became non-infectious when exposed to a temperature of 133 degrees C for 90 minutes (Duan 2003).
 - UV radiation for 60 minutes also destroyed the infective capability of the virus (Duan 2003), which may indicate that the virus may perish on surfaces exposed to sunlight for an extended period of time (however it is too soon to know for certain).
 - Cases of SARS rose with pollution (the number of inhalable particles) (Xie et al 2004), which may be an analog for disease propagation in smoky conditions.
- **Protocols for dealing with infected persons, or those suspected of infection**
 - If you have traveled in a vehicle with or been within 1 meter of someone known to be infected, evaluation in a medical setting is recommended or if that is not available, then self-quarantine and monitoring symptoms for 14 days is recommended (Chinese CDC 2020).
 - When dealing with an infected person (or one who is suspected to be infected), wearing a gown, head-covering, N95 filtering facepiece respirator, eye protection, and gloves is the protocol for viruses such as Ebola; workers who did not deploy this protection initially in China were infected (Brosseau 2020).
- **Strategies to reduce disease propagation at the community level**
 - Two strategies to reduce the disease propagation include mitigation (defined as slowing the spread) and suppression (in which the number of cases is slowed to low levels, with that situation being maintained for some period of time, potentially up to 18 months) (Ferguson et al 2020).

- If a mitigation strategy is pursued, it might include a set of non-pharmaceutical interventions including placing suspected cases on home isolation and quarantining other family members, and using social distancing of all high-risk cases including the elderly (Ferguson et al 2020). Such a strategy would be likely to reduce peak healthcare demand by 2/3, based on simulation studies, but health systems including ICUs and resources such as ventilators would be overwhelmed and thousands of deaths would likely result (Ferguson et al 2020)
 - A suppression strategy in the US would require at a minimum social distancing of the whole population, accompanied by home isolation of those with COVID-19 and quarantine of family members (Ferguson et al 2020).
- As a vaccine may not be available for 18 months or more, suppression may be necessary for a long time period, with standards being relaxed at times as the disease becomes less frequent. However, simulations indicate that the disease levels would quickly rebound, making a return to a suppression strategy necessary to avoid widespread deaths. Unfortunately, a suppression strategy would carry serious economic impacts (Ferguson et al 2020).
- The effect of public health interventions such as social distancing and shelter in place on “flattening the curve”, or reducing the rate at which COVID-19 is spreading can be viewed in interactive graphs here: <https://www.nytimes.com/interactive/2020/03/13/opinion/coronavirus-trump-response.html> (Kristof and Thompson 2020). Such strategies mean that hospitals may be able to handle all cases, and buy time for a vaccine to be developed and deployed. (Chart below appeared in Roberts 2020)



- In summary, COVID-19 cases are likely to come in waves, with vaccine development and deployment possibly being 18 months away. The severity of the waves (number of cases) can be reduced by aggressive public health measures.

Disclaimer: It should be noted that the author of this document, Dr. Karin Riley, is a Research Ecologist and not an epidemiologist. As such, Dr. Riley has expertise in research and reviewing

literature in general, but no expertise in epidemiology. However, this white paper has been reviewed and fact-checked by an infectious disease epidemiologist and an individual researching the disease genomics. Because of the evolving nature of this pandemic, information that is the best available today may soon be outdated by newer studies.



Karin L. Riley, PhD
Research Ecologist

Forest Service
Fire Sciences Lab, Rocky Mountain Research Station

p: 406-329-4806 (I am currently teleworking and not checking this number regularly)

c: 406-533-5820

karin.l.riley@usda.gov

5775 W. Broadway St.

Missoula, MT 59808

www.fs.fed.us



Caring for the land and serving people

References

- Achenbach, Joel. 2020. The coronavirus isn't mutating quickly, suggesting a vaccine would offer lasting protection. Washington Post, https://www.washingtonpost.com/health/the-coronavirus-isnt-mutating-quickly-suggesting-a-vaccine-would-offer-lasting-protection/2020/03/24/406522d6-6dfd-11ea-b148-e4ce3fbd85b5_story.html, accessed March 27, 2020.
- Brosseau, Lisa. 2020. Commentary: COVID-19 transmission messages should hinge on science. Center for Infectious Disease Research and Policy, University of Minnesota, <http://www.cidrap.umn.edu/news-perspective/2020/03/commentary-covid-19-transmission-messages-should-hinge-science>, accessed March 21, 2020
- Centers for Disease Control. 2020. Morbidity and mortality weekly report: severe outcomes among patients with Coronavirus Disease 2019 (COVID-19) – United States, March 12-March 16, 2020. <https://www.cdc.gov/mmwr/volumes/69/wr/mm6912e2.htm>, accessed March 22, 2020
- Chen, Stephen. 2020. Coronavirus can travel twice as far as official “safe distance” and stay in air for 30 minutes, Chinese study finds. South China Morning Post, <https://www.scmp.com/news/china/science/article/3074351/coronavirus-can-travel-twice-far-official-safe-distance-and-stay>, accessed March 22, 2020
- Chinese Center for Disease Control and Prevention. 2020. General questions: COVID-19 prevention and control. <http://www.chinacdc.cn/en/COVID19/202002/P020200306358351973919.pdf>
- Duan, SM, XS Zhao, RF Wen, JJ Huang, GH Pi, Zhang SX, J Han, SL Bi, L Ruan, XP Dong. 2003. Stability of SARS coronavirus in human specimens and environment and its sensitivity to heating and UV irradiation. *Biomedical and Environmental Sciences* 16(3): 246-255.

- Feng, Emily. 2020. Mystery in Wuhan: recovered coronavirus patients tested negative . . .and then positive. National Public Radio, <https://www.npr.org/sections/goatsandsoda/2020/03/27/822407626/mystery-in-wuhan-recovered-coronavirus-patients-test-negative-then-positive> , accessed March 29, 2020.
- Ferguson, Neil M., Daniel Laydon, Gemma Nedjati-Gilani, Natsuko Imai, Kylie Ainslie, Marc Baguelin, Sangeeta Bhatia, Adhiratha Boonyasiri, Zulma Cucunubá, Gina Cuomo-Dannenburg, Amy Dighe, Ilaria Dorigatti, Han Fu, Katy Gaythorpe, Will Green, Arran Hamlet, Wes Hinsley, Lucy C Okell, Sabine van Elsland, Hayley Thompson, Robert Verity, Erik Volz, Haowei Wang, Yuanrong Wang, Patrick GT Walker, Caroline Walters, Peter Winskill, Charles Whittaker, Christl A Donnelly, Steven Riley, Azra C Ghani. 2020. Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. Imperial College COVID-19 Response Team, London, UK.
- Ghani, A. C., C. A. Donnelly, D. R. Cox, J. T. Griffin, C. Fraser, T. H. Lam, L. M. Ho, W. S. Chan, R. M. Anderson, A. J. Hedley, and G. M. Leung. 2005. Methods for estimating the case fatality ratio for a novel, emerging infectious disease. *American Journal of Epidemiology* 162(5): 479-486.
- Greenfieldboyce, Nell. 2020. Do you get immunity after recovering from a case of the coronavirus? National Public Radio, <https://www.npr.org/sections/goatsandsoda/2020/03/20/819038431/do-you-get-immunity-after-recovering-from-a-case-of-coronavirus> , accessed March 27, 2020.
- John, T. Jacob and Reuben Samuel. 2000. Herd immunity and herd effect: new insights and definitions. *European Journal of Epidemiology* 16: 601-606.
- Johns Hopkins University. 2020. Coronavirus COVID-19 global cases by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University. <https://systems.jhu.edu/>, accessed March 27, 2020.
- Johnson, David L., Kenneth R. Mead, Robert A. Lynch, Deborah V. L. Hirst. 2013 Lifting the lid on toilet plume aerosol: a literature review with suggestions for future research. *American Journal of Infection Control* 41: 254-258.
- Kristof, Nicholas and Stuart A. Thompson. 2020. How much worse the coronavirus could get, in charts. <https://www.nytimes.com/interactive/2020/03/13/opinion/coronavirus-trump-response.html>, accessed March 21, 2020
- Lauer, Stephen A., Kyra H. Grantz, Qifang Bi, Forrest K. Jones, Qulu Zheng, Hannah R. Meredith, Andrew S. Azman, Nicholas G. Reich, and Justin Lessler. 2020. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Annals of Internal Medicine* doi:10.7326/M20-0504
- Li, Ruiyun, Sen Pei, Bin Chen, Yimeng Song, Tao Zhang, Wan Yang, and Jeffrey Shaman. 2020. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). *Science* DOI: 10.1126/science.abb3221

- Lu, Xiaoxia, Liqiong Zhang, Hui Du, Jingjing Zhang, Yuan Y. Li, Jingyu Qu, Wenxin Zhang, Youjie Wang, Shuangshuang Bao, Ying Li, Chuansha Wu, Hongxiu Liu, Di Liu, Jianbo Shao, Xuehua Peng, Yonghong Yang, Zhisheng Liu, Yun Xiang, Furong Zhang, Rona M. Silva, Kent E. Pinkerton, Kunling Shen, Han Xiao, Shunqing Xu, Gary W.K. Wong. 2020. SARS-CoV-2 infection in children. *New England Journal of Medicine* DOI: 10.1056/NEJMc2005073
- May, Brandon. 2020. COVID-19: Risk factors identified for poor prognosis, in-hospital mortality. *Pulmonology Advisor*
<https://www.pulmonologyadvisor.com/home/topics/lung-infection/covid-19-risk-factors-identified-for-poor-prognosis-in-hospital-mortality/>, accessed March 22, 2020.
- Mehta, Puja, Daniel F McAuley, Michael Brown, Emilie Sanchez, Rachel S Tattersall, Jessica J Manson. 2020. COVID-19: consider cytokine storm syndromes and immunosuppression. *The Lancet* [https://doi.org/10.1016/50140-6736\(20\)30628-0](https://doi.org/10.1016/50140-6736(20)30628-0)
- Moncion, K., K. Young, M. Tunis, S. Rempel, R. Stirling, and L. Zhao. 2019. Effectiveness of hand hygiene practices in preventing influenza virus infection in the community setting: a systematic review. *Canada Communicable Disease Report* 45(1):12-23.
- Moriarty, Leah F., Mateusz M. Plucinski, Barbara J. Marston, Ekaterina V. Kurbatova, Barbara Knust, Erin L. Murray, Nicki Pesik, Dale Rose, David Fitter, Miwako Kobayashi, Mitsuru Toda, Paul T. Canty, Tara Scheuer, Eric S. Halsey, Nicole J. Cohen, Lauren Stockman, Debra A. Wadford, Alexandra M. Medley, Gary Green, Joanna J. Regan, Kara Tardivel, Stefanie White, Clive Brown, Christina Morales, Cynthia Yen, Beth Wittry, Amy Freeland, Sara Naramore, Ryan T. Novak, David Daigle, Michelle Weinberg, Anna Acosta, Carolyn Herzig, Bryan K Kapella, Kathleen R. Jacobson, Katherine Lamba, Atsuyoshi Ishizumi, John Sarisky, Erik Svendsen, Tricia Blocher, Christine Wu, Julia Charles, Riley Wagner, Andrea Stewart, Paul S. Mead, Elizabeth Kurylo, Stefanie Campbell, Rachel Murray, Paul Weidle, Martin Cetron, Cindy R. Friedman, CDC Cruise Ship Response Team, California Department of Public Health COVID-19 Team, Solano County COVID-19 Team. 2020. Public Health Responses to COVID-19 Outbreaks on Cruise Ships — Worldwide, February–March 2020.
https://www.cdc.gov/mmwr/volumes/69/wr/mm6912e3.htm?s_cid=mm6912e3_w, accessed March 28, 2020/.
- Prakash, Nidhi and Ellie Hall. 2020. Doctors and nurses say more people are dying of COVID-19 in the US than we know. *Buzzfeed News*,
<https://www.buzzfeednews.com/article/nidhiprakash/coronavirus-update-dead-covid-19-doctors-hospitals>, accessed March 27, 2020.
- Regan, Helen. 2020. How can the coronavirus spread through bathroom pipes? Experts are investigating in Hong Kong. <https://www.cnn.com/2020/02/12/asia/hong-kong-coronavirus-pipes-intl-hnk/index.html>, accessed March 23, 2020

- Roberts, Siobhan. 2020. Flattening the Coronavirus curve. New York Times, <https://www.nytimes.com/2020/03/11/science/coronavirus-curve-mitigation-infection.html>, accessed March 21, 2020.
- ScienceDaily. 2020. New coronavirus stable for hours on surfaces. <https://www.sciencedaily.com/releases/2020/03/200317150116.htm>, accessed March 21, 2020
- Tang, Xiaolu, Changcheng Wu, Xiang Li, Yuhe Song, Xinmin Yao, Xinkai Wu, Yuange Duan, Hong Zhang, Yirong Wang, Zhaohui Qian, Jie Cui, and Jian Lu. In Press. On the origin and continuing evolution of SARS-CoV-2. National Science Review nwaa036, <https://doi.org/10.1093/nsr/nwaa036>
- Van Doremalen, Neeltje, Dylan H. Morris, Myndi G. Holbrook, Amandine Gamble, Brandi N. Williamson, Azaibi Tamin, Jennifer L. Harcourt, Natalie J. Thornburg, Susan I. Gerber, James O. Lloyd-Smith, Emmie de Wit, Vincent J. Munster. 2020. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. New England Journal of Medicine DOI: 10.1056/NEJMc2004973
- Weiss, Paul and David R. Murdoch. 2020. Clinical course and mortality risk of severe COVID-19. The Lancet [https://doi.org/10.1016/S0140-6736\(20\)30633-4](https://doi.org/10.1016/S0140-6736(20)30633-4)
- Wölfel, Roman, Victor M. Corman, Wolfgang Guggemos, Michael Seilmaier, Sabine Zange, Marcel A. Müller, Daniela Niemeyer, Terence C. Jones Kelly, Patrick Vollmar, Camilla Rothe, Michael Hoelscher, Tobias Bleicker, Sebastian Brünink, Julia Schneider, Rosina Ehmann, Katrin Zwirgmaier, Christian Drosten, Clemens Wendtner. 2020. Virological assessment of hospitalized cases of coronavirus disease 2019. medRxiv preprint <https://www.medrxiv.org/content/10.1101/2020.03.05.20030502v1.full.pdf>
- World Health Organization. 2020. Coronavirus disease (COVID-19) Situation Report-59, March 19, 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200319-sitrep-59-covid-19.pdf?sfvrsn=c3dcdef9_2, accessed March 22, 2020.
- Xie, Mei, Tao Lui, Xiao-an Lu, Zu-zhao Huang, and Gui-xiong Liang. 2004. Correlation research between SARS' propagation incidence and concentration of inhalable particulates in air in Guangzhou. Journal of Safety and Environment http://en.cnki.com.cn/Article_en/CJFDTotal-AQHJ200403019.htm, accessed March 22, 2020